

**TECHNOLOGY** Some graphing utilities, such as *Derive*, *Maple*, *Mathcad*, *Mathematica*, and the *TI-89*, perform symbolic differentiation. Others perform *numerical differentiation* by finding values of derivatives using the formula

$$f'(x) \approx \frac{f(x + \Delta x) - f(x - \Delta x)}{2\Delta x}$$

where  $\Delta x$  is a small number such as 0.001. Can you see any problems with this definition? For instance, using this definition, what is the value of the derivative of  $f(x) = |x|$  when  $x = 0$ ?

### THEOREM 2.1 Differentiability Implies Continuity

If  $f$  is differentiable at  $x = c$ , then  $f$  is continuous at  $x = c$ .

**Proof** You can prove that  $f$  is continuous at  $x = c$  by showing that  $f(x)$  approaches  $f(c)$  as  $x \rightarrow c$ . To do this, use the differentiability of  $f$  at  $x = c$  and consider the following limit.

$$\begin{aligned} \lim_{x \rightarrow c} [f(x) - f(c)] &= \lim_{x \rightarrow c} \left[ (x - c) \left( \frac{f(x) - f(c)}{x - c} \right) \right] \\ &= \left[ \lim_{x \rightarrow c} (x - c) \right] \left[ \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \right] \\ &= (0)[f'(c)] \\ &= 0 \end{aligned}$$



Because the difference  $f(x) - f(c)$  approaches zero as  $x \rightarrow c$ , you can conclude that  $\lim_{x \rightarrow c} f(x) = f(c)$ . So,  $f$  is continuous at  $x = c$ .  $\square$

You can summarize the relationship between continuity and differentiability as follows.


1. If a function is differentiable at  $x = c$ , then it is continuous at  $x = c$ . So, differentiability implies continuity.
2. It is possible for a function to be continuous at  $x = c$  and not be differentiable at  $x = c$ . So, continuity does not imply differentiability.

## EXERCISES FOR SECTION 2.1

In Exercises 1 and 2, estimate the slope of the graph at the point  $(x, y)$ .



In Exercises 3 and 4, use the graph shown in the figure. To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).



3. Identify or sketch each of the quantities on the figure.

(a)  $f(1)$  and  $f(4)$                       (b)  $f(4) - f(1)$

(c)  $y = \frac{f(4) - f(1)}{4 - 1}(x - 1) + f(1)$

4. Insert the proper inequality symbol ( $<$  or  $>$ ) between the given quantities.

(a)  $\frac{f(4) - f(1)}{4 - 1}$    $\frac{f(4) - f(3)}{4 - 3}$

(b)  $\frac{f(4) - f(1)}{4 - 1}$    $f'(1)$

In Exercises 5–10, find the slope of the tangent line to the graph of the function at the specified point.

5.  $f(x) = 3 - 2x$ ,  $(-1, 5)$
6.  $g(x) = \frac{3}{2}x + 1$ ,  $(-2, -2)$
7.  $g(x) = x^2 - 4$ ,  $(1, -3)$
8.  $g(x) = 5 - x^2$ ,  $(2, 1)$
9.  $f(t) = 3t - t^2$ ,  $(0, 0)$
10.  $h(t) = t^2 + 3$ ,  $(-2, 7)$

In Exercises 11–24, find the derivative by the limit process.

11.  $f(x) = 3$
12.  $g(x) = -5$
13.  $f(x) = -5x$
14.  $f(x) = 3x + 2$
15.  $h(s) = 3 + \frac{2}{3}s$
16.  $f(x) = 9 - \frac{1}{2}x$
17.  $f(x) = 2x^2 + x - 1$
18.  $f(x) = 1 - x^2$
19.  $f(x) = x^3 - 12x$
20.  $f(x) = x^3 + x^2$
21.  $f(x) = \frac{1}{x-1}$
22.  $f(x) = \frac{1}{x^2}$
23.  $f(x) = \sqrt{x+1}$
24.  $f(x) = \frac{4}{\sqrt{x}}$

In Exercises 25–32, (a) find an equation of the tangent line to the graph of  $f$  at the indicated point, (b) use a graphing utility to graph the function and its tangent line at the point, and (c) use the derivative feature of a graphing utility to confirm your results.

25.  $f(x) = x^2 + 1$ ,  $(2, 5)$
26.  $f(x) = x^2 + 2x + 1$ ,  $(-3, 4)$
27.  $f(x) = x^3$ ,  $(2, 8)$
28.  $f(x) = x^3 + 1$ ,  $(1, 2)$
29.  $f(x) = \sqrt{x}$ ,  $(1, 1)$
30.  $f(x) = \sqrt{x-1}$ ,  $(5, 2)$
31.  $f(x) = x + \frac{4}{x}$ ,  $(4, 5)$
32.  $f(x) = \frac{1}{x+1}$ ,  $(0, 1)$

In Exercises 33–36, find an equation of the line that is tangent to the graph of  $f$  and parallel to the given line.


Function	Line
33. $f(x) = x^3$	$3x - y + 1 = 0$
34. $f(x) = x^3 + 2$	$3x - y - 4 = 0$
35. $f(x) = \frac{1}{\sqrt{x}}$	$x + 2y - 6 = 0$
36. $f(x) = \frac{1}{\sqrt{x-1}}$	$x + 2y + 7 = 0$

37. The tangent line to the graph of  $y = g(x)$  at the point  $(5, 2)$  passes through the point  $(9, 0)$ . Find  $g(5)$  and  $g'(5)$ .
38. The tangent line to the graph of  $y = h(x)$  at the point  $(-1, 4)$  passes through the point  $(3, 6)$ . Find  $h(-1)$  and  $h'(-1)$ .


### Getting at the Concept

In Exercises 39–42, the graph of  $f$  is given. Select the graph of  $f'$ .


39.




40.




41.




42.




(a)




(b)



(c)



(d)



43. Sketch a graph of a function whose derivative is always negative.
44. Sketch a graph of a function whose derivative is always positive.
45. Assume that  $f'(c) = 3$ . Find  $f'(-c)$  if (a)  $f$  is an odd function and if (b)  $f$  is an even function.
46. Determine whether the limit yields the derivative of a differentiable function  $f$ . Explain.

(a)  $\lim_{\Delta x \rightarrow 0} \frac{f(x + 2\Delta x) - f(x)}{2\Delta x}$


(b)  $\lim_{\Delta x \rightarrow 0} \frac{f(x + 2) - f(x)}{\Delta x}$

(c)  $\lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x - \Delta x)}{2\Delta x}$


(d)  $\lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}$

In Exercises 47 and 48, find equations of the two tangent lines to the graph of  $f$  that pass through the indicated point.


47.  $f(x) = 4x - x^2$



48.  $f(x) = x^2$



49. **Graphical Reasoning** The figure shows the graph of  $g'$ .



- (a)  $g'(0) =$
- (b)  $g'(3) =$
- (c) What can you conclude about the graph of  $g$  knowing that  $g'(1) = -\frac{8}{3}$ ?
- (d) What can you conclude about the graph of  $g$  knowing that  $g'(-4) = \frac{7}{3}$ ?
- (e) Is  $g(6) - g(4)$  positive or negative? Explain.
- (f) Is it possible to find  $g(2)$  from the graph? Explain.

50. **Graphical Reasoning** Use a graphing utility to graph each function and its tangent lines when  $x = -1$ ,  $x = 0$ , and  $x = 1$ . Based on the results, determine whether the slope of a tangent line to the graph of a function is always distinct for different values of  $x$ .

(a)  $f(x) = x^2$       (b)  $g(x) = x^3$

51. **Graphical, Numerical, and Analytic Analysis** In Exercises 51 and 52, use a graphing utility to graph  $f$  on the interval  $[-2, 2]$ . Complete the table by graphically estimating the slopes of the graph at the indicated points. Then evaluate the slopes analytically and compare your results with those obtained graphically.

$x$	-2	-1.5	-1	-0.5	0	0.5	1	1.5	2
$f(x)$									
$f'(x)$									

51.  $f(x) = \frac{1}{4}x^3$

52.  $f(x) = \frac{1}{2}x^2$

53. **Graphical Reasoning** In Exercises 53 and 54, use a graphing utility to graph the functions  $f$  and  $g$  in the same viewing window where

$$g(x) = \frac{f(x + 0.01) - f(x)}{0.01}$$

Label the graphs and describe the relationship between them.

53.  $f(x) = 2x - x^2$

54.  $f(x) = 3\sqrt{x}$

In Exercises 55 and 56, evaluate  $f(2)$  and  $f'(2)$  and use the results to approximate  $f'(2)$ .

55.  $f(x) = x(4 - x)$

56.  $f(x) = \frac{1}{4}x^3$

57. **Graphical Reasoning** In Exercises 57 and 58, use a graphing utility to graph the function and its derivative in the same viewing window. Label the graphs and describe the relationship between them.

57.  $f(x) = \frac{1}{\sqrt{x}}$

58.  $f(x) = \frac{x^3}{4} - 3x$

59. **Writing** In Exercises 59 and 60, consider the functions  $f$  and  $S_{\Delta x}$  where

$$S_{\Delta x}(x) = \frac{f(2 + \Delta x) - f(2)}{\Delta x}(x - 2) + f(2)$$

- (a) Use a graphing utility to graph  $f$  and  $S_{\Delta x}$  in the same viewing window for  $\Delta x = 1, 0.5$ , and  $0.1$ .
- (b) Give a written description of the graphs of  $S$  for the different values of  $\Delta x$  in part (a).

59.  $f(x) = 4 - (x - 3)^2$

60.  $f(x) = x + \frac{1}{x}$

In Exercises 61–70, use the alternative form of the derivative to find the derivative at  $x = c$  (if it exists).

61.  $f(x) = x^2 - 1$ ,  $c = 2$

62.  $g(x) = x(x - 1)$ ,  $c = 1$

63.  $f(x) = x^3 + 2x^2 + 1$ ,  $c = -2$

64.  $f(x) = x^3 + 2x$ ,  $c = 1$

65.  $g(x) = \sqrt{|x|}$ ,  $c = 0$

66.  $f(x) = 1/x$ ,  $c = 3$

67.  $f(x) = (x - 6)^{2/3}$ ,  $c = 6$

68.  $g(x) = (x + 3)^{1/3}$ ,  $c = -3$



69.  $h(x) = |x + 5|$ ,  $c = -5$

70.  $f(x) = |x - 4|$ ,  $c = 4$


In Exercises 71–80, describe the  $x$ -values at which  $f$  is differentiable.

71.  $f(x) = |x + 3|$


72.  $f(x) = |x^2 - 9|$




73.  $f(x) = \frac{1}{x+1}$




75.  $f(x) = (x-3)^{2/3}$




77.  $f(x) = \sqrt{x-1}$




79.  $f(x) = \begin{cases} 4 - x^2, & x > 0 \\ x^2 - 4, & x \leq 0 \end{cases}$




80.  $f(x) = \begin{cases} x^2 - 2x, & x > 1 \\ x^3 - 3x^2 + 3x, & x \leq 1 \end{cases}$




74.  $f(x) = \frac{2x}{x-1}$



76.  $f(x) = x^{2/5}$



78.  $f(x) = \frac{x^2}{x^2 - 4}$



In Exercises 81–84, find the derivatives from the left and from the right at  $x = 1$  (if they exist). Is the function differentiable at  $x = 1$ ?

81.  $f(x) = |x - 1|$

82.  $f(x) = \sqrt{1 - x^2}$


83.  $f(x) = \begin{cases} (x-1)^3, & x \leq 1 \\ (x-1)^2, & x > 1 \end{cases}$

84.  $f(x) = \begin{cases} x, & x \leq 1 \\ x^2, & x > 1 \end{cases}$

In Exercises 85 and 86, determine whether the function is differentiable at  $x = 2$ .

85.  $f(x) = \begin{cases} x^2 + 1, & x \leq 2 \\ 4x - 3, & x > 2 \end{cases}$

86.  $f(x) = \begin{cases} \frac{1}{2}x + 1, & x < 2 \\ \sqrt{2x}, & x \geq 2 \end{cases}$

 **87. Graphical Reasoning** A line with slope  $m$  passes through the point  $(0, 4)$  and has the equation  $y = mx + 4$ .

(a) Write the distance  $d$  between the line and the point  $(3, 1)$  as a function of  $m$ .

(b) Use a graphing utility to graph the function  $d$  in part (a). Based on the graph, is the function differentiable at every value of  $m$ ? If not, where is it not differentiable?

**88. Conjecture** Consider the functions  $f(x) = x^2$  and  $g(x) = x^3$ .

(a) Graph  $f$  and  $f'$  on the same set of axes.

(b) Graph  $g$  and  $g'$  on the same set of axes.

(c) Identify any pattern between the functions  $f$  and  $g$  and their respective derivatives. Use the pattern to make a conjecture about  $h'(x)$  if  $h(x) = x^n$ , where  $n$  is an integer and  $n \geq 2$ .

(d) Find  $f'(x)$  if  $f(x) = x^4$ . Compare the result with the conjecture in part (c). Is this a proof of your conjecture? Explain.

**True or False?** In Exercises 89–92, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

**89.** The slope of the tangent line to the differentiable function  $f$  at the point  $(2, f(2))$  is

$$\frac{f(x + \Delta x) - f(x)}{\Delta x}$$


**90.** If a function is continuous at a point, then it is differentiable at that point.

**91.** If a function has derivatives from both the right and the left at a point, then it is differentiable at that point.

**92.** If a function is differentiable at a point, then it is continuous at that point.

**93.** Let  $f(x) = \begin{cases} x \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$  and  $g(x) = \begin{cases} x^2 \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$


Show that  $f$  is continuous, but not differentiable, at  $x = 0$ . Show that  $g$  is differentiable at 0, and find  $g'(0)$ .

 **94. Writing** Use a graphing utility to graph the two functions  $f(x) = x^2 + 1$  and  $g(x) = |x| + 1$  in the same viewing window. Use the *zoom* and *trace* features to analyze the graphs near the point  $(0, 1)$ . What do you observe? Which function is differentiable at this point? Write a short paragraph describing the geometric significance of differentiability at a point.


## EXERCISES FOR SECTION 2.2

In Exercises 1 and 2, use the graph to estimate the slope of the tangent line to  $y = x^n$  at the point  $(1, 1)$ . Verify your answer analytically. To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).


1. (a)  $y = x^{1/2}$




(b)  $y = x^{3/2}$




(c)  $y = x^2$




(d)  $y = x^3$




2. (a)  $y = x^{-1/2}$




(b)  $y = x^{-1}$



(c)  $y = x^{-3/2}$



(d)  $y = x^{-2}$



In Exercises 3–24, find the derivative of the function.

3.  $y = 8$

4.  $f(x) = -2$

5.  $y = x^6$

6.  $y = x^8$

7.  $y = \frac{1}{x^7}$

8.  $y = \frac{1}{x^8}$

9.  $f(x) = \sqrt[3]{x}$

10.  $g(x) = \sqrt[4]{x}$

11.  $f(x) = x + 1$

12.  $g(x) = 3x - 1$

13.  $f(t) = -2t^2 + 3t - 6$

14.  $y = t^2 + 2t - 3$

15.  $g(x) = x^2 + 4x^3$

16.  $y = 8 - x^3$

17.  $s(t) = t^3 - 2t + 4$

18.  $f(x) = 2x^3 - x^2 + 3x$

19.  $y = \frac{\pi}{2} \sin \theta - \cos \theta$

20.  $g(t) = \pi \cos t$

21.  $y = x^2 - \frac{1}{2} \cos x$

22.  $y = 5 + \sin x$

23.  $y = \frac{1}{x} - 3 \sin x$

24.  $y = \frac{5}{(2x)^3} + 2 \cos x$

In Exercises 25–30, complete the table, using Example 6 as a model.

	Original Function	Rewrite	Differentiate	Simplify
25.	$y = \frac{5}{2x^2}$			
26.	$y = \frac{2}{3x^2}$			
27.	$y = \frac{3}{(2x)^3}$			
28.	$y = \frac{\pi}{(3x)^2}$			
29.	$y = \frac{\sqrt{x}}{x}$			
30.	$y = \frac{4}{x^{-3}}$			

In Exercises 31–38, find the slope of the graph of the function at the indicated point. Use the *derivative* feature of a graphing utility to confirm your results.

Function	Point
31. $f(x) = \frac{3}{x^2}$	(1, 3)
32. $f(t) = 3 - \frac{3}{5t}$	$(\frac{3}{5}, 2)$
33. $f(x) = -\frac{1}{2} + \frac{7}{5}x^3$	$(0, -\frac{1}{2})$
34. $y = 3x^3 - 6$	(2, 18)
35. $y = (2x + 1)^2$	(0, 1)
36. $f(x) = 3(5 - x)^2$	(5, 0)
37. $f(\theta) = 4 \sin \theta - \theta$	(0, 0)
38. $g(t) = 2 + 3 \cos t$	$(\pi, -1)$

$-16t^2 + 16t^{-1} = 0$   
 $-16(t^2 + t^{-1}) = 0$   
 $-16(t^{-1}(t^3 + 1)) = 0$

In Exercises 39–52, find the derivative of the function.

39.  $f(x) = x^2 + 5 - 3x^{-2}$

40.  $f(x) = x^2 - 3x - 3x^{-2}$

41.  $g(t) = t^2 - \frac{4}{t^3}$

42.  $f(x) = x + \frac{1}{x^2}$

43.  $f(x) = \frac{x^3 - 3x^2 + 4}{x^2}$

44.  $h(x) = \frac{2x^2 - 3x + 1}{x}$

45.  $y = x(x^2 + 1)$

46.  $y = 3x(6x - 5x^2)$

47.  $f(x) = \sqrt{x} - 6\sqrt[3]{x}$

48.  $f(x) = \sqrt[3]{x} + \sqrt{x}$

49.  $h(s) = s^{4/5} - s^{2/3}$

50.  $f(t) = t^{2/3} - t^{1/3} + 4$

51.  $f(x) = 6\sqrt{x} + 5 \cos x$

52.  $f(x) = \frac{2}{\sqrt[3]{x}} + 3 \cos x$

**In Exercises 53–56, (a) find an equation of the tangent line to the graph of  $f$  at the indicated point, (b) use a graphing utility to graph the function and its tangent line at the point, and (c) use the derivative feature of a graphing utility to confirm your results.**

Function	Point
53. $y = x^4 - 3x^2 + 2$	(1, 0)
54. $y = x^3 + x$	(-1, -2)
55. $f(x) = \frac{2}{4\sqrt{x^3}}$	(1, 2)
56. $y = (x^2 + 2x)(x + 1)$	(1, 6)

**In Exercises 57–62, determine the point(s) (if any) at which the graph of the function has a horizontal tangent line.**


57.  $y = x^4 - 8x^2 + 2$       58.  $y = x^3 + x$   
 59.  $y = \frac{1}{x^2}$       60.  $y = x^2 + 1$   
 61.  $y = x + \sin x, 0 \leq x < 2\pi$   
 62.  $y = \sqrt{3}x + 2 \cos x, 0 \leq x < 2\pi$

**In Exercises 63–66, find  $k$  such that the line is tangent to the graph of the function.**

Function	Line
63. $f(x) = x^2 - kx$	$y = 4x - 9$
64. $f(x) = k - x^2$	$y = -4x + 7$
65. $f(x) = \frac{k}{x}$	$y = -\frac{3}{4}x + 3$
66. $f(x) = k\sqrt{x}$	$y = x + 4$

### Getting at the Concept

67. Use the graph of  $f$  to answer each question. To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).





- (a) Between which two consecutive points is the average rate of change of the function greatest?  
 (b) Is the average rate of change of the function between A and B greater than or less than the instantaneous rate of change at B?  
 (c) Sketch a tangent line to the graph between C and D such that the slope of the tangent line is the same as the average rate of change of the function between C and D.
68. Sketch the graph of a function  $f$  such that  $f' > 0$  for all  $x$  and the rate of change of the function is decreasing.

### Getting at the Concept (continued)

**In Exercises 69 and 70, the relationship between  $f$  and  $g$  is given. Give the relationship between  $f'$  and  $g'$ .**

69.  $g(x) = f(x) + 6$       70.  $g(x) = -5f(x)$

**In Exercises 71 and 72, the graphs of a function  $f$  and its derivative  $f'$  are shown on the same set of coordinate axes. Label the graphs as  $f$  or  $f'$  and write a short paragraph stating the criteria used in making the selection. To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).**



73. Sketch the graphs of  $y = x^2$  and  $y = -x^2 + 6x - 5$ , and sketch the two lines that are tangent to both graphs. Find equations of these lines.  
 74. Show that the graphs of the two equations  $y = x$  and  $y = 1/x$  have tangent lines that are perpendicular to each other at their point of intersection.

**In Exercises 75 and 76, find an equation of the tangent line to the graph of the function  $f$  through the point  $(x_0, y_0)$  not on the graph. To find the point of tangency  $(x, y)$  on the graph of  $f$ , solve the equation**

$$f'(x) = \frac{y_0 - y}{x_0 - x}$$

75.  $f(x) = \sqrt{x}$       76.  $f(x) = \frac{2}{x}$   
 $(x_0, y_0) = (-4, 0)$        $(x_0, y_0) = (5, 0)$

77. **Linear Approximation** Use a graphing utility (in square mode) to zoom in on the graph of  $f(x) = 4 - \frac{1}{2}x^2$  to approximate  $f'(1)$ . Use the derivative to find  $f'(1)$ .  
 78. **Linear Approximation** Use a graphing utility (in square mode) to zoom in on the graph of  $f(x) = 4\sqrt{x} + 1$  to approximate  $f'(4)$ . Use the derivative to find  $f'(4)$ .  
 79. **Linear Approximation** Consider the function  $f(x) = x^{3/2}$  with the solution point  $(4, 8)$ .  
 (a) Use a graphing utility to obtain the graph of  $f$ . Use the zoom feature to obtain successive magnifications of the graph in the neighborhood of the point  $(4, 8)$ . After zooming in a few times, the graph should appear nearly linear. Use the trace feature to determine the coordinates of a point “near”  $(4, 8)$ . Find an equation of the secant line  $S(x)$  through the two points.

- (b) Find the equation of the line

$$T(x) = f'(4)(x - 4) + f(4)$$

tangent to the graph of  $f$  passing through the given point. Why are the linear functions  $S$  and  $T$  nearly the same?

- (c) Use a graphing utility to graph  $f$  and  $T$  on the same set of coordinate axes. Note that  $T$  is a "good" approximation of  $f$  when  $x$  is "close to" 4. What happens to the accuracy of the approximation as you move farther away from the point of tangency?
- (d) Demonstrate the conclusion in part (c) by completing the table.

$\Delta x$	-3	-2	-1	-0.5	-0.1	0
$f(4 + \Delta x)$						
$T(4 + \Delta x)$						

$\Delta x$	0.1	0.5	1	2	3
$f(4 + \Delta x)$					
$T(4 + \Delta x)$					

80. **Linear Approximation** Repeat Exercise 79 for the function  $f(x) = x^3$  where  $T(x)$  is the line tangent to the graph at the point  $(1, 1)$ . Explain why the accuracy of the linear approximation decreases more rapidly than in Exercise 79.

**True or False?** In Exercises 81–86, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

81. If  $f'(x) = g'(x)$ , then  $f(x) = g(x)$ .  
 82. If  $f(x) = g(x) + c$ , then  $f'(x) = g'(x)$ .  
 83. If  $y = \pi^2$ , then  $dy/dx = 2\pi$ .  
 84. If  $y = x/\pi$ , then  $dy/dx = 1/\pi$ .  
 85. If  $g(x) = 3f(x)$ , then  $g'(x) = 3f'(x)$ .  
 86. If  $f(x) = 1/x^n$ , then  $f'(x) = 1/(nx^{n-1})$ .

In Exercises 87–90, find the average rate of change of the function over the indicated interval. Compare this average rate of change with the instantaneous rates of change at the endpoints of the interval.

Function	Interval
87. $f(t) = 2t + 7$	$[1, 2]$
88. $f(t) = t^2 - 3$	$[2, 2.1]$
89. $f(x) = \frac{-1}{x}$	$[1, 2]$
90. $f(x) = \sin x$	$\left[0, \frac{\pi}{6}\right]$



**Vertical Motion** In Exercises 91 and 92, use the position function  $s(t) = -16t^2 + v_0t + s_0$  for free-falling objects.

91. A silver dollar is dropped from the top of a building that is 1362 feet tall.
- Determine the position and velocity functions for the coin.
  - Determine the average velocity on the interval  $[1, 2]$ .
  - Find the instantaneous velocities when  $t = 1$  and  $t = 2$ .
  - Find the time required for the coin to reach ground level.
  - Find the velocity of the coin at impact.
92. A ball is thrown straight down from the top of a 220-foot building with an initial velocity of  $-22$  feet per second. What is its velocity after 3 seconds? What is its velocity after falling 108 feet?



**Vertical Motion** In Exercises 93 and 94, use the position function  $s(t) = -4.9t^2 + v_0t + s_0$  for free-falling objects.

93. A projectile is shot upward from the surface of earth with an initial velocity of 120 meters per second. What is its velocity after 5 seconds? After 10 seconds?
94. To estimate the height of a building, a stone is dropped from the top of the building into a pool of water at ground level. How high is the building if the splash is seen 6.8 seconds after the stone is dropped?


**Think About It** In Exercises 95 and 96, the graph of a position function is shown. It represents the distance in miles that a person drives during a 10-minute trip to work. Make a sketch of the corresponding velocity function.



**Think About It** In Exercises 97 and 98, the graph of a velocity function is shown. It represents the velocity in miles per hour during a 10-minute drive to work. Make a sketch of the corresponding position function.



- 99. Modeling Data** The stopping distance of an automobile traveling at a speed  $v$  (kilometers per hour) is the distance  $R$  (meters) the car travels during the reaction time of the driver plus the distance  $B$  (meters) the car travels after the brakes are applied (see figure). The table shows the results of an experiment.



$v$	20	40	60	80	100
$R$	3.3	6.7	10.0	13.3	16.7
$B$	2.3	8.9	20.2	35.9	56.7

- Use the regression capabilities of a graphing utility to find a linear model for reaction time.
  - Use the regression capabilities of a graphing utility to find a quadratic model for braking time.
  - Determine the polynomial giving the total stopping distance  $T$ .
  - Use a graphing utility to graph the functions  $R$ ,  $B$ , and  $T$  in the same viewing window.
  - Find the derivative of  $T$  and the rate of change of the total stopping distance for  $v = 40$ ,  $v = 80$ , and  $v = 100$ .
  - Use the results of this exercise to draw conclusions about the total stopping distance as speed increases.
- 100. Velocity** Verify that the average velocity over the time interval  $[t_0 - \Delta t, t_0 + \Delta t]$  is the same as the instantaneous velocity at  $t = t_0$  for the position function
- $$s(t) = -\frac{1}{2}at^2 + c.$$
- 101. Area** The area of a square with sides of length  $s$  is given by  $A = s^2$ . Find the rate of change of the area with respect to  $s$  when  $s = 4$  meters.
- 102. Volume** The volume of a cube with sides of length  $s$  is given by  $V = s^3$ . Find the rate of change of the volume with respect to  $s$  when  $s = 4$  centimeters.
- 103. Inventory Management** The annual inventory cost  $C$  for a certain manufacturer is

$$C = \frac{1,008,000}{Q} + 6.3Q$$

where  $Q$  is the order size when the inventory is replenished. Find the change in annual cost when  $Q$  is increased from 350 to 351, and compare this with the instantaneous rate of change when  $Q = 350$ .

- 104. Fuel Cost** A car is driven 15,000 miles a year and gets  $x$  miles per gallon. Assume that the average fuel cost is \$1.25 per gallon. Find the annual cost of fuel  $C$  as a function of  $x$  and use this function to complete the table.

$x$	10	15	20	25	30	35	40
$C$							
$\frac{dC}{dx}$							

Who would benefit more from a 1-mile-per-gallon increase in fuel efficiency—the driver of a car that gets 15 miles per gallon or the driver of a car that gets 35 miles per gallon? Explain.

- 105. Writing** The number of gallons  $N$  of regular unleaded gasoline sold by a gasoline station at a price of  $p$  dollars per gallon is given by  $N = f(p)$ .
- Describe the meaning of  $f'(1.479)$ .
  - Is  $f'(1.479)$  usually positive or negative? Explain.
- 106. Newton's Law of Cooling** This law states that the rate of change of the temperature of an object is proportional to the difference between the object's temperature  $T$  and the temperature  $T_a$  of the surrounding medium. Write an equation for this law.
- 107.** Find an equation of the parabola  $y = ax^2 + bx + c$  that passes through  $(0, 1)$  and is tangent to the line  $y = x - 1$  at  $(1, 0)$ .
- 108.** Let  $(a, b)$  be an arbitrary point on the graph of  $y = 1/x$ ,  $x > 0$ . Prove that the area of the triangle formed by the tangent line through  $(a, b)$  and the coordinate axes is 2.
- 109.** Find the tangent line(s) to the curve  $y = x^3 - 9x$  through the point  $(1, -9)$ .
- 110.** Find the equation(s) of the tangent line(s) to the parabola  $y = x^2$  through the given point.
- $(0, a)$
  - $(a, 0)$
- Are there any restrictions on the constant  $a$ ?
- 111.** Find  $a$  and  $b$  such that
- $$f(x) = \begin{cases} ax^3, & x \leq 2 \\ x^2 + b, & x > 2 \end{cases}$$
- is differentiable everywhere.
- 112.** Where are the functions  $f_1(x) = |\sin x|$  and  $f_2(x) = \sin |x|$  differentiable?
- 113.** Prove that  $\frac{d}{dx} [\cos x] = -\sin x$ .

**FOR FURTHER INFORMATION** For a geometric interpretation of the derivatives of trigonometric functions, see the article "Sines and Cosines of the Times" by Victor J. Katz in *Math Horizons*. To view this article, go to the website [www.matharticles.com](http://www.matharticles.com).



## EXERCISES FOR SECTION 2.3

In Exercises 1–6, use the Product Rule to differentiate the function.

1.  $g(x) = (x^2 + 1)(x^2 - 2x)$

2.  $f(x) = (6x + 5)(x^3 - 2)$

3.  $h(t) = \sqrt[3]{t}(t^2 + 4)$

4.  $g(s) = \sqrt{s}(4 - s^2)$

5.  $f(x) = x^3 \cos x$

6.  $g(x) = \sqrt{x} \sin x$

In Exercises 7–12, use the Quotient Rule to differentiate the function.

7.  $f(x) = \frac{x}{x^2 + 1}$

8.  $g(t) = \frac{t^2 + 2}{2t - 7}$

9.  $h(x) = \frac{\sqrt[3]{x}}{x^3 + 1}$

10.  $h(s) = \frac{s}{\sqrt{s} - 1}$

11.  $g(x) = \frac{\sin x}{x^2}$

12.  $f(t) = \frac{\cos t}{t^3}$

In Exercises 13–18, find  $f'(x)$  and  $f''(c)$ .

Function	Value of $c$
13. $f(x) = (x^3 - 3x)(2x^2 + 3x + 5)$	$c = 0$
14. $f(x) = (x^2 - 2x + 1)(x^3 - 1)$	$c = 1$
15. $f(x) = \frac{x^2 - 4}{x - 3}$	$c = 1$
16. $f(x) = \frac{x + 1}{x - 1}$	$c = 2$
17. $f(x) = x \cos x$	$c = \frac{\pi}{4}$
18. $f(x) = \frac{\sin x}{x}$	$c = \frac{\pi}{6}$

In Exercises 19–24, complete the table without using the Quotient Rule (see Example 6).

Function	Rewrite	Differentiate	Simplify
19. $y = \frac{x^2 + 2x}{3}$			
20. $y = \frac{5x^2 - 3}{4}$			
21. $y = \frac{7}{3x^3}$			
22. $y = \frac{4}{5x^2}$			
23. $y = \frac{4x^{3/2}}{x}$			
24. $y = \frac{3x^2 - 5}{7}$			

In Exercises 25–38, find the derivative of the algebraic function.

25.  $f(x) = \frac{3\sqrt{-2x - x^2}}{x^2 - 1}$

26.  $f(x) = \frac{x^3 + 3x + 2}{x^2 - 1}$

27.  $f(x) = x\left(1 - \frac{4}{x+3}\right)$

28.  $f(x) = x^4\left(1 - \frac{2}{x+1}\right)$

29.  $f(x) = \frac{2x + 5}{\sqrt{x}}$

30.  $f(x) = \sqrt[3]{x}(\sqrt{x} + 3)$

31.  $h(s) = (s^3 - 2)^2$

32.  $h(x) = (x^2 - 1)^2$

33.  $f(x) = \frac{2 - \frac{1}{x}}{x - 3}$

34.  $g(x) = x^2\left(\frac{2}{x} - \frac{1}{x+1}\right)$

35.  $f(x) = (3x^3 + 4x)(x - 5)(x + 1)$

36.  $f(x) = (x^2 - x)(x^2 + 1)(x^2 + x + 1)$

37.  $f(x) = \frac{x^2 + c^2}{x^2 - c^2}$ ,  $c$  is a constant

38.  $f(x) = \frac{c^2 - x^2}{c^2 + x^2}$ ,  $c$  is a constant

In Exercises 39–54, find the derivative of the trigonometric function.

39.  $f(t) = t^2 \sin t$

40.  $f(\theta) = (\theta + 1) \cos \theta$

41.  $f(t) = \frac{\cos t}{t}$

42.  $f(x) = \frac{\sin x}{x}$

43.  $f(x) = -x + \tan x$

44.  $y = x + \cot x$

45.  $g(t) = \sqrt[4]{t} + 8 \sec t$

46.  $h(s) = \frac{1}{s} - 10 \csc s$

47.  $y = \frac{3(1 - \sin x)}{2 \cos x}$

48.  $y = \frac{\sec x}{x}$

49.  $y = -\csc x - \sin x$

50.  $y = x \sin x + \cos x$

51.  $f(x) = x^2 \tan x$

52.  $f(x) = \sin x \cos x$

53.  $y = 2x \sin x + x^2 \cos x$

54.  $h(\theta) = 5\theta \sec \theta + \theta \tan \theta$

In Exercises 55–58, use a computer algebra system to differentiate the function.

55.  $g(x) = \left(\frac{x+1}{x+2}\right)(2x-5)$

56.  $f(x) = \left(\frac{x^2 - x - 3}{x^2 + 1}\right)(x^2 + x + 1)$

57.  $g(\theta) = \frac{\theta}{1 - \sin \theta}$

58.  $f(\theta) = \frac{\sin \theta}{1 - \cos \theta}$

In Exercises 59–62, evaluate the derivative of the function at the indicated point. Use a graphing utility to verify your result.

Function	Point
59. $y = \frac{1 + \csc x}{1 - \csc x}$	$\left(\frac{\pi}{6}, -3\right)$
60. $f(x) = \tan x \cot x$	$(1, 1)$
61. $h(t) = \frac{\sec t}{t}$	$\left(\pi, -\frac{1}{\pi}\right)$
62. $f(x) = \sin x(\sin x + \cos x)$	$\left(\frac{\pi}{4}, 1\right)$

In Exercises 63–68, (a) find an equation of the tangent line to the graph of  $f$  at the indicated point, (b) use a graphing utility to graph the function and its tangent line at the point, and (c) use the derivative feature of a graphing utility to confirm your results.

Function	Point
63. $f(x) = (x^3 - 3x + 1)(x + 2)$	(1, -3)
64. $f(x) = (x - 1)(x^2 - 2)$	(0, 2)
65. $f(x) = \frac{x}{x - 1}$	(2, 2)
66. $f(x) = \frac{(x - 1)}{(x + 1)}$	$(2, \frac{1}{3})$
67. $f(x) = \tan x$	$(\frac{\pi}{4}, 1)$
68. $f(x) = \sec x$	$(\frac{\pi}{3}, 2)$

In Exercises 69 and 70, determine the point(s) at which the graph of the function has a horizontal tangent.

69.  $f(x) = \frac{x^2}{x - 1}$       70.  $f(x) = \frac{x^2}{x^2 + 1}$

In Exercises 71 and 72, verify that  $f'(x) = g'(x)$ , and explain the relationship between  $f$  and  $g$ .

71.  $f(x) = \frac{3x}{x + 2}$ ,  $g(x) = \frac{5x + 4}{x + 2}$   
 72.  $f(x) = \frac{\sin x - 3x}{x}$ ,  $g(x) = \frac{\sin x + 2x}{x}$

In Exercises 73 and 74, find the derivative of the function  $f$  for  $n = 1, 2, 3$ , and 4. Use the result to write a general rule for  $f'(x)$  in terms of  $n$ .

73.  $f(x) = x^n \sin x$       74.  $f(x) = \frac{\cos x}{x^n}$

75. **Area** The length of a rectangle is given by  $2t + 1$  and its height is  $\sqrt{t}$ , where  $t$  is time in seconds and the dimensions are in centimeters. Find the rate of change of the area with respect to time.

76. **Volume** The radius of a right circular cylinder is given by  $\sqrt{t + 2}$  and its height is  $\frac{1}{2}\sqrt{t}$ , where  $t$  is time in seconds and the dimensions are in inches. Find the rate of change of the volume with respect to time.

77. **Inventory Replenishment** The ordering and transportation cost  $C$  for the components used in manufacturing a certain product is

$$C = 100\left(\frac{200}{x^2} + \frac{x}{x + 30}\right), \quad x \geq 1$$

where  $C$  is measured in thousands of dollars and  $x$  is the order size in hundreds. Find the rate of change of  $C$  with respect to  $x$  when (a)  $x = 10$ , (b)  $x = 15$ , and (c)  $x = 20$ . What do these rates of change imply about increasing order size?

78. **Boyle's Law** This law states that if the temperature of a gas remains constant, its pressure is inversely proportional to its volume. Use the derivative to show that the rate of change of the pressure is inversely proportional to the square of the volume.

79. **Population Growth** A population of 500 bacteria is introduced into a culture and grows in number according to the equation

$$P(t) = 500\left(1 + \frac{4t}{50 + t^2}\right)$$

where  $t$  is measured in hours. Find the rate at which the population is growing when  $t = 2$ .

80. **Rate of Change** Determine whether there exist any values of  $x$  in the interval  $[0, 2\pi)$  such that the rate of change of  $f(x) = \sec x$  and the rate of change of  $g(x) = \csc x$  are equal.

81. Prove the following differentiation rules.

(a)  $\frac{d}{dx}[\sec x] = \sec x \tan x$

(b)  $\frac{d}{dx}[\csc x] = -\csc x \cot x$

(c)  $\frac{d}{dx}[\cot x] = -\csc^2 x$

82. **Modeling Data** The table shows the number of motor homes  $n$  (in thousands) in the United States and the retail value  $v$  (in millions of dollars) of these motor homes for the years 1992 through 1997. The year is represented by  $t$ , with  $t = 2$  corresponding to 1992. (Source: Recreation Vehicle Industry Association)

Year	1992	1993	1994	1995	1996	1997
$n$	226.3	243.8	306.7	281.0	274.6	239.3
$v$	\$6963	\$7544	\$9897	\$9768	\$9788	\$9139

(a) Use a graphing utility to find quadratic models for the number of motor homes  $n(t)$  and the total retail value  $v(t)$  of the motor homes.

(b) Find  $A = v(t)/n(t)$ . What does this function represent?

(c) Find  $A'(t)$ . Interpret the derivative in the context of these data.

In Exercises 83–88, find the second derivative of the function.

83.  $f(x) = 4x^{3/2}$

84.  $f(x) = x + 32x^{-2}$

85.  $f(x) = \frac{x}{x - 1}$

86.  $f(x) = \frac{x^2 + 2x - 1}{x}$

87.  $f(x) = 3 \sin x$

88.  $f(x) = \sec x$

In Exercises 89–92, find the higher-order derivative.

Given

Find

89.  $f'(x) = x^2$

$f''(x)$

90.  $f''(x) = 2 - \frac{2}{x}$

$f'''(x)$

91.  $f'''(x) = 2\sqrt{x}$

$f^{(4)}(x)$

92.  $f^{(4)}(x) = 2x + 1$

$f^{(6)}(x)$

**Getting at the Concept**

93. Sketch the graph of a differentiable function  $f$  such that  $f(2) = 0$ ,  $f' < 0$  for  $-\infty < x < 2$ , and  $f' > 0$  for  $2 < x < \infty$ .
94. Sketch the graph of a differentiable function  $f$  such that  $f > 0$  and  $f' < 0$  for all real numbers  $x$ .

In Exercises 95–98, find  $f'(2)$  given the following.


$g(2) = 3$  and  $g'(2) = -2$

$h(2) = -1$  and  $h'(2) = 4$

95.  $f(x) = 2g(x) + h(x)$       96.  $f(x) = 4 - h(x)$

97.  $f(x) = \frac{g(x)}{h(x)}$       98.  $f(x) = g(x)h(x)$

In Exercises 99 and 100, the graphs of  $f, f'$ , and  $f''$  are shown on the same set of coordinate axes. Which is which? To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).



101. **Acceleration** The velocity of an object in meters per second is  $v(t) = 36 - t^2$ ,  $0 \leq t \leq 6$ .

Find the velocity and acceleration of the object when  $t = 3$ . What can be said about the speed of the object when the velocity and acceleration have opposite signs?

102. **Stopping Distance** A car is traveling at a rate of 66 feet per second (45 miles per hour) when the brakes are applied. The position function for the car is

$s(t) = -8.25t^2 + 66t$

where  $s$  is measured in feet and  $t$  is measured in seconds. Use this function to complete the table, and find the average velocity during each time interval.

$t$	0	1	2	3	4
$s(t)$					
$v(t)$					
$a(t)$					

103. **Acceleration** An automobile's velocity starting from rest is

$v(t) = \frac{100t}{2t + 15}$

where  $v$  is measured in feet per second. Find the acceleration at each of the following times.

- (a) 5 seconds    (b) 10 seconds    (c) 20 seconds

104. **Finding a Pattern** Develop a general rule for  $f^{(n)}(x)$  if

(a)  $f(x) = x^n$     and    (b)  $f(x) = \frac{1}{x}$ .

105. **Finding a Pattern** Consider the function  $f(x) = g(x)h(x)$ .

(a) Use the product rule to generate rules for finding  $f''(x)$ ,  $f'''(x)$ , and  $f^{(4)}(x)$ .

(b) Use the results in part (a) to write a general rule for  $f^{(n)}(x)$ .

106. **Finding a Pattern** Develop a general rule for  $[xf(x)]^{(n)}$  where  $f$  is a differentiable function of  $x$ .

**Linear and Quadratic Approximations** The linear and quadratic approximations of a function  $f$  at  $x = a$  are

$P_1(x) = f'(a)(x - a) + f(a)$  and

$P_2(x) = \frac{1}{2}f''(a)(x - a)^2 + f'(a)(x - a) + f(a)$ .

In Exercises 107 and 108, (a) find the specified linear and quadratic approximations of  $f$ , (b) use a graphing utility to graph  $f$  and the approximations, (c) determine whether  $P_1$  or  $P_2$  is the better approximation, and (d) state how the accuracy changes as you move farther from  $x = a$ .

107.  $f(x) = \cos x$

108.  $f(x) = \sin x$

$a = \frac{\pi}{3}$

$a = \frac{\pi}{2}$

**True or False?** In Exercises 109–114, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

109. If  $y = f(x)g(x)$ , then  $dy/dx = f'(x)g'(x)$ .
110. If  $y = (x + 1)(x + 2)(x + 3)(x + 4)$ , then  $d^5y/dx^5 = 0$ .
111. If  $f'(c)$  and  $g'(c)$  are zero and  $h(x) = f(x)g(x)$ , then  $h'(c) = 0$ .
112. If  $f(x)$  is an  $n$ th-degree polynomial, then  $f^{(n+1)}(x) = 0$ .
113. The second derivative represents the rate of change of the first derivative.
114. If the velocity of an object is constant, then its acceleration is zero.

115. Find the derivative of  $f(x) = x|x|$ . Does  $f''(0)$  exist?

116. **Think About It** Let  $f$  and  $g$  be functions whose first and second derivatives exist on an interval  $I$ . Which of the following formulas is (are) true?

(a)  $fg'' - f''g = (fg' - f'g)'$

(b)  $fg'' + f''g = (fg)''$

## REVIEW EXERCISES FOR CHAPTER 2

**2.1** In Exercises 1–4, find the derivative of the function by using the definition of the derivative.

1.  $f(x) = x^2 - 2x + 3$

2.  $f(x) = \frac{x+1}{x-1}$



3.  $f(x) = \sqrt{x} + 1$

4.  $f(x) = \frac{2}{x}$

In Exercises 5 and 6, describe the  $x$ -values at which  $f$  is differentiable.

5.  $f(x) = (x+1)^{2/3}$

6.  $f(x) = \frac{4x}{x+3}$



7. Sketch the graph of  $f(x) = 4 - |x - 2|$ .

(a) Is  $f$  continuous at  $x = 2$ ?

(b) Is  $f$  differentiable at  $x = 2$ ? Explain.

8. Sketch the graph of  $f(x) = \begin{cases} x^2 + 4x + 2, & x < -2 \\ 1 - 4x - x^2, & x \geq -2 \end{cases}$

(a) Is  $f$  continuous at  $x = -2$ ?

(b) Is  $f$  differentiable at  $x = -2$ ? Explain.

In Exercises 9 and 10, find the slope of the tangent line to the graph of the function at the specified point.

9.  $g(x) = \frac{2}{3}x^2 - \frac{x}{6}$ ,  $\left(-1, \frac{5}{6}\right)$

10.  $h(x) = \frac{3x}{8} - 2x^2$ ,  $\left(-2, -\frac{35}{4}\right)$

**2.2** In Exercises 11 and 12, (a) find an equation of the tangent line to the graph of  $f$  at the indicated point, (b) use a graphing utility to graph the function and its tangent line at the point, and (c) use the derivative feature of the graphing utility to confirm your results.

11.  $f(x) = x^3 - 1$ ,  $(-1, -2)$



12.  $f(x) = \frac{2}{x+1}$ ,  $(0, 2)$

In Exercises 13 and 14, use the alternative form of the derivative to find the derivative at  $x = c$  (if it exists).

13.  $g(x) = x^2(x-1)$ ,  $c = 2$

14.  $f(x) = \frac{1}{x+1}$ ,  $c = 2$

**Writing** In Exercises 15 and 16, the figure shows the graphs of a function and its derivative. Label the graphs as  $f$  or  $f'$  and write a short paragraph stating the criteria used in making the selection. To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).



**2.2** In Exercises 17–32, find the derivative of the function.

17.  $y = 25$

18.  $y = -12$

19.  $f(x) = x^8$

20.  $g(x) = x^{12}$

21.  $h(t) = 3t^4$

22.  $f(t) = -8t^5$

23.  $f(x) = x^3 - 3x^2$

24.  $g(s) = 4s^4 - 5s^2$

25.  $h(x) = 6\sqrt{x} + 3\sqrt[3]{x}$

26.  $f(x) = x^{1/2} - x^{-1/2}$

27.  $g(t) = \frac{2}{3t^2}$

28.  $h(x) = \frac{2}{(3x)^2}$

29.  $f(\theta) = 2\theta - 3\sin\theta$

30.  $g(\alpha) = 4\cos\alpha + 6$

31.  $f(\theta) = 3\cos\theta - \frac{\sin\theta}{4}$

32.  $g(\alpha) = \frac{5\sin\alpha}{3} - 2\alpha$

33. **Vibrating String** When a guitar string is plucked, it vibrates with a frequency of  $F = 200\sqrt{T}$ , where  $F$  is measured in vibrations per second and the tension  $T$  is measured in pounds. Find the rate of change of  $F$  when (a)  $T = 4$  and (b)  $T = 9$ .

34. **Vertical Motion** A ball is dropped from a height of 100 feet. One second later, another ball is dropped from a height of 75 feet. Which ball hits the ground first?

35. **Vertical Motion** To estimate the height of a building, a weight is dropped from the top of the building into a pool at ground level. How high is the building if the splash is seen 9.2 seconds after the weight is dropped?

36. **Vertical Motion** A bomb is dropped from an airplane at an altitude of 14,400 feet. How long will it take for the bomb to reach the ground? (Because of the motion of the plane, the fall will not be vertical, but the time will be the same as that for a vertical fall.) The plane is moving at 600 miles per hour. How far will the bomb move horizontally after it is released from the plane?

**37. Projectile Motion** A ball thrown follows a path described by  $y = x - 0.02x^2$ .

- Sketch a graph of the path.
- Find the total horizontal distance the ball was thrown.
- At what  $x$ -value does the ball reach its maximum height? (Use the symmetry of the path.)
- Find an equation that gives the instantaneous rate of change of the height of the ball with respect to the horizontal change. Evaluate the equation at  $x = 0, 10, 25, 30$ , and  $50$ .
- What is the instantaneous rate of change of the height when the ball reaches its maximum height?

**38. Projectile Motion** The path of a projectile thrown at an angle of  $45^\circ$  with level ground is

$$y = x - \frac{32}{v_0^2}(x^2)$$

where the initial velocity is  $v_0$  feet per second.

- Find the  $x$ -coordinate of the point where the projectile strikes the ground. Use the symmetry of the path of the projectile to locate the  $x$ -coordinate of the point where the projectile reaches its maximum height.
- What is the instantaneous rate of change of the height when the projectile is at its maximum height?
- Show that doubling the initial velocity of the projectile multiplies both the maximum height and the range by a factor of 4.
- Find the maximum height and range of a projectile thrown with an initial velocity of 70 feet per second. Use a graphing utility to sketch the path of the projectile.

**39. Horizontal Motion** The position function of a particle moving along the  $x$ -axis is

$$x(t) = t^2 - 3t + 2$$

for  $-\infty < t < \infty$ .

- Find the velocity of the particle.
- Find the open  $t$ -interval(s) in which the particle is moving to the left.
- Find the position of the particle when the velocity is 0.
- Find the speed of the particle when the position is 0.

**40. Modeling Data** The speed of a car in miles per hour and the stopping distance in feet are recorded in the table.

Speed ( $x$ )	20	30	40	50	60
Stopping Distance ( $y$ )	25	55	105	188	300

- Use the regression capabilities of a graphing utility to find a quadratic model for the data.
- Use a graphing utility to plot the data and graph the model.
- Use a graphing utility to graph  $dy/dx$ .
- Use the model to approximate the stopping distance at a speed of 65 miles per hour.
- Use the graphs in parts (b) and (c) to explain the change in stopping distance as the speed increases.

**2.3** In Exercises 41–57, find the derivative of the function.

41.  $f(x) = (3x^2 + 7)(x^2 - 2x + 3)$

42.  $g(x) = (x^3 - 3x)(x + 2)$

43.  $h(x) = \sqrt{x} \sin x$

45.  $f(x) = \frac{2x^3 - 1}{x^2}$

47.  $f(x) = \frac{x^2 + x - 1}{x^2 - 1}$

49.  $f(x) = \frac{1}{4 - 3x^2}$

51.  $y = \frac{x^2}{\cos x}$

53.  $y = 3x^2 \sec x$

55.  $y = -x \tan x$

57.  $y = x \cos x - \sin x$

44.  $f(t) = t^3 \cos t$

46.  $f(x) = \frac{x + 1}{x - 1}$

48.  $f(x) = \frac{6x - 5}{x^2 + 1}$

50.  $f(x) = \frac{9}{3x^2 - 2x}$

52.  $y = \frac{\sin x}{x^2}$

54.  $y = 2x - x^2 \tan x$

56.  $y = \frac{1 + \sin x}{1 - \sin x}$

**58. Acceleration** The velocity of an object in meters per second is  $v(t) = 36 - t^2$ ,  $0 \leq t \leq 6$ . Find the velocity and acceleration of the object when  $t = 4$ .

In Exercises 59–62, find the second derivative of the function.

59.  $g(t) = t^3 - 3t + 2$

60.  $f(x) = 12\sqrt[4]{x}$

61.  $f(\theta) = 3 \tan \theta$

62.  $h(t) = 4 \sin t - 5 \cos t$

In Exercises 63 and 64, show that the function satisfies the equation.

Function  $y = 2 \sin x + 3 \cos x$

Equation  $y'' + y = 0$

63.  $y = 2 \sin x + 3 \cos x$

$y'' + y = 0$

64.  $y = \frac{10 - \cos x}{x}$

$xy' + y = \sin x$

**2.4** In Exercises 65–80, find the derivative of the function.

65.  $f(x) = \sqrt{1 - x^3}$

66.  $f(x) = \sqrt[3]{x^2 - 1}$

67.  $h(x) = \left(\frac{x - 3}{x^2 + 1}\right)^2$

68.  $f(x) = \left(x^2 + \frac{1}{x}\right)^5$

69.  $f(s) = (s^2 - 1)^{5/2}(s^3 + 5)$

70.  $h(\theta) = \frac{\theta}{(1 - \theta)^3}$

71.  $y = 3 \cos(3x + 1)$

72.  $y = 1 - \cos 2x + 2 \cos^2 x$

73.  $y = \frac{1}{2} \csc 2x$

74.  $y = \csc 3x + \cot 3x$

75.  $y = \frac{x}{2} - \frac{\sin 2x}{4}$

76.  $y = \frac{\sec^7 x}{7} - \frac{\sec^5 x}{5}$

77.  $y = \frac{2}{3} \sin^{3/2} x - \frac{2}{7} \sin^{7/2} x$

78.  $f(x) = \frac{3x}{\sqrt{x^2 + 1}}$

79.  $y = \frac{\sin \pi x}{x + 2}$

80.  $y = \frac{\cos(x - 1)}{x - 1}$

In Exercises 81–88, use a computer algebra system to find the derivative of the function. Use the utility to graph the function and its derivative on the same set of coordinate axes. Describe the behavior of the function that corresponds to any zeros of the graph of the derivative.

81.  $f(t) = t^2(t - 1)^5$       82.  $f(x) = [(x - 2)(x + 4)]^2$   
 83.  $g(x) = \frac{2x}{\sqrt{x + 1}}$       84.  $g(x) = x\sqrt{x^2 + 1}$   
 85.  $f(t) = \sqrt{t + 1} \sqrt[3]{t + 1}$       86.  $y = \sqrt{3x(x + 2)^3}$   
 87.  $y = \tan\sqrt{1 - x}$       88.  $y = 2 \csc^3(\sqrt{x})$

In Exercises 89–92, find the second derivative of the function.

89.  $y = 2x^2 + \sin 2x$       90.  $y = \frac{1}{x} + \tan x$   
 91.  $f(x) = \cot x$       92.  $y = \sin^2 x$

In Exercises 93–96, use a computer algebra system to find the second derivative of the function.

93.  $f(t) = \frac{t}{(1 - t)^2}$       94.  $g(x) = \frac{6x - 5}{x^2 + 1}$   
 95.  $g(\theta) = \tan 3\theta - \sin(\theta - 1)$       96.  $h(x) = x\sqrt{x^2 - 1}$

97. **Refrigeration** The temperature  $T$  of food put in a freezer is

$$T = \frac{700}{t^2 + 4t + 10}$$

where  $t$  is the time in hours. Find the rate of change of  $T$  with respect to  $t$  at each of the following times.

- (a)  $t = 1$     (b)  $t = 3$     (c)  $t = 5$     (d)  $t = 10$

98. **Fluid Flow** The emergent velocity  $v$  of a liquid flowing from a hole in the bottom of a tank is given by  $v = \sqrt{2gh}$ , where  $g$  is the acceleration due to gravity (32 feet per second per second) and  $h$  is the depth of the liquid in the tank. Find the rate of change of  $v$  with respect to  $h$  when (a)  $h = 9$  and (b)  $h = 4$ . (Note that  $g = +32$  feet per second per second. The sign of  $g$  depends on how a problem is modeled. In this case, letting  $g$  be negative would produce an imaginary value for  $v$ .)

**2.5** In Exercises 99–104, use implicit differentiation to find  $dy/dx$ .

99.  $x^2 + 3xy + y^3 = 10$       100.  $x^2 + 9y^2 - 4x + 3y = 0$   
 101.  $y\sqrt{x} - x\sqrt{y} = 16$       102.  $y^2 = (x - y)(x^2 + y)$   
 103.  $x \sin y = y \cos x$       104.  $\cos(x + y) = x$

In Exercises 105 and 106, find the equations of the tangent line and the normal line to the graph of the equation at the indicated point. Use a graphing utility to graph the equation, the tangent line, and the normal line.

105.  $x^2 + y^2 = 20$ , (2, 4)      106.  $x^2 - y^2 = 16$ , (5, 3)

**2.6**


107. A point moves along the curve  $y = \sqrt{x}$  in such a way that the  $y$ -value is increasing at a rate of 2 units per second. At what rate is  $x$  changing for each of the following values?

- (a)  $x = \frac{1}{2}$     (b)  $x = 1$     (c)  $x = 4$

108. **Surface Area** The edges of a cube are expanding at a rate of 5 centimeters per second. How fast is the surface area changing when each edge is 4.5 centimeters?

109. **Changing Depth** The cross section of a 5-meter trough is an isosceles trapezoid with a 2-meter lower base, a 3-meter upper base, and an altitude of 2 meters. Water is running into the trough at a rate of 1 cubic meter per minute. How fast is the water level rising when the water is 1 meter deep?

110. **Linear and Angular Velocity** A rotating beacon is located 1 kilometer off a straight shoreline (see figure). If the beacon rotates at a rate of 3 revolutions per minute, how fast (in kilometers per hour) does the beam of light appear to be moving to a viewer who is  $\frac{1}{2}$  kilometer down the shoreline?



Not drawn to scale

111. **Moving Shadow** A sandbag is dropped from a balloon at a height of 60 meters when the angle of elevation to the sun is  $30^\circ$  (see figure). Find the rate at which the shadow of the sandbag is traveling along the ground when the sandbag is at a height of 35 meters. [Hint: The position of the sandbag is given by  $s(t) = 60 - 4.9t^2$ .]

